

Application No. 10/023,265
Docket No. 2000U057.US
Reply to Office Action Dated May 06, 2004

Amendments to the Claims

page 1, before the first paragraph:

--CROSS-REFERENCE TO RELATED APPLICATION

The present application claims priority to Provisional U.S. Application U.S.S.N. 60/258,482 filed on December 28, 2000, which is herein incorporated by reference.--

page 2, first paragraph:

(Amended) For US purposes the following references are mentioned: US 4,845,067; US 4,999,327; JP 1126111; US 4,508,842; and UK 1015054.

Application No. 10/023,265
Docket No. 2000U057.US
Reply to Office Action Dated May 06, 2004


Amendments to the Claims

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1-13 (Cancelled)


14. (Currently amended) An A polymerization catalyst comprising a combination of at least one activator and a reaction product of a transition metal compound with a tridentate ligand generating composition represented by a formula of:



(I)

or


Application No. 10/023,265
 Docket No. 2000U057.US
 Reply to Office Action Dated May 06, 2004



wherein: R₂ and R₃ are hydrocarbyl radicals or substituted hydrocarbyl radicals, R₅ - R₈ are each, independently, hydrogen, a hydrocarbyl radical or a substituted hydrocarbyl radical; one of R₁, R₂, R₃, R₄, or R₉ is a radical that contains a Group 16 atom and R* is a hydrocarbyl radical or substituted hydrocarbyl radical when R₁ is a radical that contains a Group 16 atom, otherwise R₁, R₂, R₃, R₄, R₉ and R* are each, independently, hydrogen, a hydrocarbyl radical or a substituted hydrocarbyl radical; and for formula (I) m and n are values of 0 or 1, and when m is 0 and n is 0 R₂ and R₃ may be joined together to form an aromatic ring structure, and when n is 0 and m is 1 R₂ and R₃ may be joined together to form ring structures; any two adjacent groups of R₅ to R₉ may be joined together to form ring structures; for formula (II) R₁ through R₉ and R* are as explained above and R₁₀ is hydrogen, a hydrocarbyl radical or a substituted hydrocarbyl radical; and p, q and r are values of 0 or 1 wherein p is 0 only when q is 1 and r is 0.


Application No. 10/023,265
Docket No. 2000U057.US
Reply to Office Action Dated May 06, 2004

15. (Original) The polymerization catalyst of claim 14 wherein the tridentate ligand generating compound is represented by the formula:



wherein R_4 is a radical that contains an oxygen based functional group selected from an alcohol, an aldehyde, a ketone, or an epoxide and R_5 and R_9 are alkyl radicals.

16. (Currently amended) The polymerization catalyst of claim 14 wherein the tridentate ligand generating compound is represented by the formula:




Application No. 10/023,265

Docket No. 2000U057.US

Reply to Office Action Dated May 06, 2004

wherein R_1 is a radical that contains an oxygen based functional group ~~such as an alcohol, an aldehyde, a ketone, an epoxide~~ and R^* , R_4 , R_5 , R_9 , and R_{10} are hydrocarbyl radicals.

17. (Original) The polymerization catalyst of claim 14 wherein the tridentate ligand generating compound is represented by the formula:



wherein R_1 is a radical that contains an oxygen based functional group selected from an alcohol, an aldehyde, a ketone, an epoxide and R^* , R_2 , R_4 , R_5 , R_7 , R_9 , and R_{10} are hydrocarbyl radicals.

18. (Original) The polymerization catalyst of claim 14 wherein the transition metal compound is of a Group 4 metal.
19. (Original) The polymerization catalyst of claim 18 wherein the transition metal is Zr.

Application No. 10/023,265
Docket No. 2000U057.US
Reply to Office Action Dated May 06, 2004

20. (Currently amended) The polymerization catalyst of claim 14 wherein the ~~oxygen containing ligand of the catalyst~~ Group 16 atom, when bonded to the transition metal, forms a ring of 5 to 8 atoms.
21. (Currently amended) The polymerization catalyst of claim 14 wherein the ~~oxygen containing ligand of the catalyst~~ Group 16 atom, when bonded to the transition metal, forms a ring of 5 to 7 atoms.
22. (Currently amended) The polymerization catalyst of claim 14 wherein the ~~oxygen containing ligand of the catalyst~~ Group 16 atom, when bonded to the transition metal, forms a ring of 6 atoms.
23. (Currently amended) The polymerization catalyst of claim 14 wherein the ~~oxygen based functional group radical that contains a Group 16 atom~~ is a ketone.
24. (Currently amended) The polymerization catalyst of claim 14 wherein the ~~oxygen based functional group radical that contains a Group 16 atom~~ is an alcohol.
25. (Original) The polymerization catalyst of claim 14 wherein the a Group 16 atom is a sulfur based functional group.
- 26-27 (Cancelled)